



2023 Lower Long Lake Water Quality Review

Introduction

The goals of this testing protocol were to monitor various water quality parameters of the lake, compare results to historical data, and identify any potential risks to the health of Lower Long Lake. Water samples were taken at four different locations and tested for 14 parameters. Tests were conducted once in the spring and once in the late summer. Tests were conducted with a YSI ProDSS Multiparameter Water Quality Meter or LaMotte SMART2 Colorimeter.

Test results were compared to historical data from the report “2022 Lower Long Lake Water Quality Review” by LakePro, Inc.

In this report, we included historical data from Water Quality Investigators. Their report provided annual averages for many of the parameters from 2002 to 2009. Including this data allows us to see more accurate trends in the water quality data. In order to make the analysis easier, we added annual averages for our data and trendlines on the graphs. The trend lines allow us to see the direction each water quality parameter is moving.

Results

Parameter	2023 Season		Status
	Average	Target Range	
Temperature	65.7 °F	Less Than 75 °F	● Healthy
Dissolved Oxygen	7.8 mg/L	4.0 – 12.0 mg/L	● Healthy
Total Phosphorus	74 ppb	0 – 100 ppb	● Healthy
Phosphate	29 ppb	0 – 100 ppb	● Healthy
Nitrate	48 ppb	0 – 200 ppb	● Healthy
Chlorophyll-a	1.7 ppb	0 – 7.3 ppb	● Healthy
Transparency	17.2 feet	More than 6.5 feet	● Healthy
pH	8.5 S.U.	7.0 – 9.0 S.U.	● Healthy
Total Dissolved Solids	504 ppm	0 – 1,000 ppm	● Healthy
Conductivity	724 ppm	0 – 1,500 ppm	● Healthy
Alkalinity	128 ppm	100 – 250 ppm	● Healthy
Sulfate	11.1 ppm	3 – 30 ppm	● Healthy
Fluoride	0.08 ppm	0.01 – 0.30 ppm	● Healthy
Chloride	141 ppm	0 – 230 ppm	● Healthy





Preface

2023 was LakePro's thirteenth year testing water quality on Lower Long Lake. The historical data reveals trends over the testing history. The trend lines on the following graphs show the change from 2002 to 2023. Each successive year of testing will provide more insight into how the lake changed.

Discussion

The results of this year's testing indicate that the water of Lower Long Lake remained outstanding throughout the summer. The results show that the aquatic environment was very suitable to support natural wildlife. Also, the lake was safe for recreational uses, such as swimming, boating, fishing, etc., as there are no signs of pollution.

The **Temperature** of the surface water was warmer than usual to start the season due to the early thaw we observed. The temperature then increased slightly into the late summer as expected. The **Dissolved Oxygen** in the lake remained excellent throughout the summer. Furthermore, the sufficient oxygen concentration was a positive attribute headed into winter when ice seals the lake off from atmospheric oxygen.

The concentrations of both **Total Phosphorus** and Phosphate were within the target range throughout the summer. **Phosphate** is the usable form of phosphorus.

The **Nitrate** concentrations remained comfortably within the target range throughout the entire summer. Although the amount was lessened, it is important that residents take measures to ensure their property is not contributing excess fertilizers to the lake.

We tested water samples for **Chlorophyll** as a direct indicator of plant production. The chlorophyll concentrations responded strongly to the water temperatures, increasing over the course of the summer and overall, less than that of the previous year. During all tests the plant pigment was within the target range at all four sites.

The **Transparency** was excellent during every test, measuring deeper than most other lakes that we tested. Transparency can be affected by many different factors, including suspended solids, dissolved solids, acids, and algae growth. The clear water is generally a positive attribute, but it also allows more sunlight to reach the lake bottom to fuel plant growth.

The **pH** was within the target during all testing events.

The **Total Dissolved Solids** and **Conductivity** consistently remained within their designated target ranges during all tests. As the summer progressed and rainfall increased, these parameters experienced a gradual rise due to heightened runoff, introducing foreign substances into the lake. Overall, this year's test shows a gradual decrease in both areas.


The **Alkalinity** was within the target range all summer but decreased throughout the summer. The major reason for the decrease in the late summer was productivity. As lake organisms become more active in the summertime, they produce more carbon dioxide. As this gas dissolves, it needs to be buffered, using up the carbonate ions. Rainfall in July infiltrated the ground and reached limestone bedrock. As this additional water reached the lake, it replenished the alkalinity in August.

The **Sulfate**, **Fluoride**, and **Chloride** were all within their target ranges for the entire summer.






Historical Trends



Temperature (°F)		
	May	August
Site 1	53.6	78.3
Site 2	54.2	78.1
Site 3	51.7	78.1
Site 4	53.2	78.5
Season Average		65.7

Target Range: < 75°F

The historical temperature trend has shown minimal alteration. Temperature fluctuations were influenced by the chosen testing dates and the annual weather variations. As we continue gathering data in the years ahead, the trend line is expected to evolve into a more precise gauge of the lake's changes.




Dissolved Oxygen		
	May	August
Site 1	8.2	7.5
Site 2	8.5	7.1
Site 3	8.3	7.2
Site 4	8.3	7.4
Season Average		7.8

Target Range: 4.0 – 12.0 mg/L

The dissolved oxygen trend showed a slight decrease over the testing history and an upward tick during last year and this year's testing events. Oxygen concentrations remained extremely healthy, showing the lake carries a healthy oxygen concentration despite temperature changes. Monitoring this trend will persist, and if a more pronounced decline occurs, we will advise additional measures accordingly.






Total Phosphorus		
	May	August
Site 1	90	70
Site 2	90	60
Site 3	80	50
Site 4	100	50
Season Average		74

Target Range: 0 – 100 ppb

The total phosphorus annual averages showed a steady increase up until 2019. The last 4 years however have showed a decrease, on par with 2022 data. Lakes generally accumulate substances, including nutrients, in the process of eutrophication. The state law banning phosphorus fertilizers and active plant management will help decrease phosphorus continuing in the future. The phosphorus concentrations were exceptionally good for a large, heavily developed watershed.




Phosphate		
	May	August
Site 1	40	30
Site 2	40	20
Site 3	30	20
Site 4	40	10
Season Average		29

Target Range: 0 – 100 ppb

Phosphate is the form of phosphorus that is usable to aquatic plants. The phosphate data also showed a slightly upward trend since we began testing this parameter in 2011. Despite the increase, the phosphate remained within the target range with a slight decline and helped to limit plant and algae growth in the lake.






Nitrate		
	May	August
Site 1	70	20
Site 2	70	20
Site 3	90	30
Site 4	60	20
Season Average		48

Target Range: 0 – 200 ppb

Nitrate is another vital nutrient for the growth of aquatic plants. Water Quality Investigators obtained high results early in the testing history, so the recent data resulted in a downward historical trend. It is important residents continue to be conscience of their property and practices to ensure more nutrients do not enter the lake.




Chlorophyll		
	May	August
Site 1	0.8	2.9
Site 2	0.6	3.0
Site 3	0.7	2.7
Site 4	0.9	2.1
Season Average		1.7

Target Range: 0 – 7.2 ppb

Chlorophyll concentrations were tested as an indicator of plant production, primarily algae in the water column. Over the testing history, the results increased steadily, which matched the increase in total phosphorus. This is also common with invasive plants in the lake, such as Eurasian Milfoil. This reinforces the urgency for responsible land management in the watershed to prevent additional phosphorus from entering the lake. Over the past 3 years however, both chlorophyll and total phosphorus have been on the lower end.






Transparency		
	May	August
Site 1	17.3	17.7
Site 2	16.9	17.3
Site 3	17.1	16.9
Site 4	17.3	17.1
Season Average		17.2

Target Range: > 6.5 feet

Transparency was affected by total dissolved solids, total suspended solids, algae growth, and rain frequency and amount. Overall, the transparency of the lake increased over the testing history



pH		
	May	August
Site 1	8.7	8.2
Site 2	8.9	8.4
Site 3	8.7	8.1
Site 4	8.8	8.1
Season Average		8.5


Target Range: 7.0 – 9.0 S.U.

There was a slight decrease in pH over the testing history, but it stayed in the target range of 7 to 9. We will look for the pH to remain level in future years. If the pH ever drops drastically, we will look for the cause of that change in order to mitigate the trend.






Target Range: 0 – 1,000 ppm



Total Dissolved Solids		
	May	August
Site 1	525	531
Site 2	521	529
Site 3	524	526
Site 4	536	540
Season Average		529

The Total Dissolved Solids showed a slight upward trend over the testing history. This is common for inland lakes as they accumulate more substances from their watershed and the atmosphere.




Conductivity		
	May	August
Site 1	718	724
Site 2	729	716
Site 3	732	720
Site 4	730	726
Season Average		724

Target Range: 0 – 1,500 μ S

Like the TDS, Conductivity showed a slight upward trend. Conductivity is an extension of TDS and measures the number of ionic molecules in the water (which conduct electricity, usually salts).






Alkalinity		
	May	August
Site 1	132	135
Site 2	135	126
Site 3	129	124
Site 4	121	119
Season Average		128

Target Range: 0 – 250 ppm

LakePro started testing this parameter in 2011. Since then, the alkalinity decreased slightly. Alkalinity works as a buffer to stabilize the pH when foreign substances enter the lake, so it is important this parameter stay at a healthy level to protect the lake from other changes.




Sulfate		
	May	August
Site 1	13.0	11.0
Site 2	16.0	10.0
Site 3	10.0	10.0
Site 4	9.0	10.0
Season Average		11.1

Target Range: 3 – 30 ppm

Calcium sulfate and magnesium sulfate are common minerals in surface water, so some sulfate should be present. Elevated levels of sulfate can indicate pollution. Over the testing history, sulfate remained within the target range and decreased toward the bottom of the target range.






Fluoride		
	May	August
Site 1	0.01	0.08
Site 2	0.04	0.11
Site 3	0.09	0.10
Site 4	0.08	0.12
Season Average		0.08

Target Range: 0.01 – 0.30 ppm

Fluoride occurs naturally in ground water, so some may be present in the lake surface water. Elevated levels can indicate pollution but are not physiologically harmful. Over the testing history, fluoride remained comfortably within the target range.



Chloride		
	May	August
Site 1	149	138
Site 2	146	135
Site 3	146	131
Site 4	137	142
Season Average		141

Target Range: 0 – 230 ppm

Chloride is a major anion found in water. This substance may be due to the natural process of water passing through salt formations in the earth or may be evidence of the intrusion of pollution from industrial processes or road salting. The trend was slightly upward due to high concentrations from 2005 to 2012. Since then, there has been a steady downward trend.





9353 Hill Road • Swartz Creek, MI 48473
(810) 635-4400 • Fax (810) 635-4404

www.lakeproinc.com

Conclusion

Overall, the water quality of Lower Long Lake was excellent again this year. All other parameters remained within their target ranges, and some were the best we saw across the state. To continue this path in keeping the waterbody healthy, it is vital that everyone within the watershed take steps to limit their nutrient input to the lake.

Despite being situated within a densely developed watershed and being bordered by residential properties, Lower Long Lake stands as a remarkable water resource boasting exceptional water quality. While there are inevitably aspects where water quality enhancements are possible, the lake consistently ranks among the top performers in our evaluations. You should take pride in your association with this lake and continue in your diligent efforts towards its ongoing enhancement.

Thanks for choosing LakePro,

Michael Smith





Analysis Information

Temperature:	The water temperature directly affects the amount of oxygen that can dissolve into the water. The temperature of surface waters is not indicative of the entire water column.
Transparency:	The ability of light to penetrate the water column is determined by the amount of dissolved and suspended particles in the water. Although aesthetically desirable, transparent water allows increased light to reach the lakebed and may result in vegetation growth.
pH:	pH is a measure of acidity or alkalinity. pH is a general measure of lake health and can roughly indicate the range of other measurements such as alkalinity and hardness.
TDS:	Total Dissolved Solids is the amount of all organic and inorganic substances in the water in a molecular or ionized state. Higher values generally indicate richer and more productive water. Lower values usually indicate cleaner and less productive water.
Conductivity:	Conductivity is a measure of the ability of water to conduct electricity. Dissolved ions in the water increase conductivity, thus TDS and Conductivity are closely related.
Alkalinity:	Alkalinity refers to the ability of the water to neutralize acids, mainly through the hydrogenation of carbonate ions. Therefore, the alkalinity is expressed as "ppm as CaCO_3 ". However, other basic molecules in the water can also contribute to alkalinity.
Dissolved Oxygen:	D.O. is a measure of the amount of oxygen dissolved in the water. This oxygen is available to fish and other animals for respiration. Vegetation generally increases DO, particularly during the day and early evening. Animals and other respiring organisms consume the oxygen, mostly during the day. Oxygen is also added to the lake through wave action, rain, fountains and aerators.
Total Phosphorus:	Phosphorus is an essential nutrient for plant growth. However, concentrations exceeding 100 ppb can impair the water and results in nuisance vegetation growth.
Phosphates:	Phosphate is the form of phosphorous that is most readily available to plants and algae.
Nitrate:	Nitrogen is also essential for plant growth. Nitrate is the predominant form of nitrogen in water. Excessive nitrate concentrations may also result in pollution and increased vegetation.
Chlorophyll-a:	Chlorophyll-a is a direct measurement of the amount of green pigment produced by plants and phytoplankton. This indicates the amount of plant growth and is used to calculate a Trophic State Index.
Sulfate:	Sulfate occurs naturally as minerals, such as calcium sulfate and magnesium sulfate. In fresh water, sulfate is usually the second or third most abundant anion. Other sources of sulfate include water material from pulp mills, steel mills, food processing operations, and municipal wastes. Under low oxygen conditions, sulfate can be reduced to hydrogen sulfide gas, which smells like rotten eggs.
Fluoride:	Fluoride may occur naturally or be added to public drinking water supplies.
Chloride:	Chloride is one of the major anions found in water and sewage. The presence of chlorides may be due to water passing through salt formations in the earth or pollution from industrial processes, domestic wastes, or road salt. The salt content of water affects the



distribution of plant and animal life in an aquatic system, based on the amount of slat they can tolerate.

Fecal Coliforms: Non-fecal coliforms are naturally found as soil organisms. Fecal Coliforms, such as *E. coli*, are coliforms found in the intestines of warm-blooded animals and humans. The presence of fecal coliforms indicates contamination from either animals or humans.

Trophic States

Oligotrophic: Water is very clear. Nutrient levels are generally low. Plant and algae productivity are also low. Sufficient dissolved oxygen in the bottom, cooler waters allow cold-water fish to survive, such as salmon and trout.

Mesotrophic: Water is moderately clear. Nutrient levels are slightly elevated. Plant and algae productivity are present, but generally not a nuisance. Oxygen and temperature in the lower portion of the lake allow walleye and perch to survive.

Eutrophic: Water is not clear due to high nutrients levels, increased turbidity, and excessive algal growth. There is no oxygen in the bottom, cooler waters, restricting the lake to warm water species, such as bass and bluegill.

Hypereutrophic: Nutrient levels are extremely high, promoting very high algae productivity. Blue-green algae blooms are likely. High turbidity and algae growth make the water opaque. Little plant growth is restricted to invasive plants. The only fish that can survive this environment are rough fish, such as carp, catfish, and mudminnows.

Sample Sites:

